



# **Stabilization of Neoclassical Tearing Mode by Electron Cyclotron Current Drive and Its Evolution Simulation on JT-60U Tokamak**

**K. Nagasaki 1), A. Isayama 2), N. Hayashi 2),  
T. Ozeki 2), M. Takechi 2), N. Oyama 2), S. Ide 2),  
S. Yamamoto 1) and JT-60 Team**

- 1) Institute of Advanced Energy, Kyoto University**
- 2) Japan Atomic Energy Research Institute**

- **Electron cyclotron current drive (ECCD) is considered to be one of attractive methods to stabilize the neoclassical tearing mode (NTM), which compensates the missing bootstrap current in the magnetic island.**
- **The necessary EC power for complete stabilization is estimated as high as 30MW in ITER. The reduction of EC power and the determination of EC conditions will help more reliable operation of high  $\beta$  plasmas without NTM.**
- **The nonlinear NTM physics is still yet understood well. It is necessary to understand the NTM physics such as the excitation and the disappearance.**

## 1. Introduction

## 2. NTM stabilization experiment by ECCD in JT-60U

Dependence of 3/2 NTM on EC injection timing

Stabilization at high  $\beta_N$  plasma using 2nd harmonic ECCD

## 3. Simulation of 3/2 NTM stabilization

Modified Rutherford equation (MRE) coupled with 1.5D transport code and EC code.



Determination of coefficients in MRE

Comparison with early and late ECCD experiment


## 4. Conclusion

# Early ECCD is more effective to suppress 3/2 NTM.


- High  $\beta_p$  ELMy H-mode plasma
- $I_p=1.5\text{MA}$ ,  $B_t=3.66\text{T}$ ,  $q_{95}=3.8$ ,  $P_{\text{NB}}=20\text{MW}$
- Only the  $m/n=3/2$  mode is excited.
- $P_{\text{EC}}=0-2.7\text{MW}$
- $I_{\text{EC}}$ : 25kA/MW, 0.17MA/m<sup>2</sup> at  $q=3/2$
- Rotation frequency: 10-12kHz



## Early ECCD






## Late ECCD







# The EC power for 3/2 NTM suppression is lower in early ECCD.

- The EC power for the complete suppression of the  $m/n=3/2$  mode is lower than in the late ECCD.
- The stabilization effect becomes weak when the current location is deviated by about half of the maximum magnetic island width.







- The  $T_e$  perturbation measured with ECE is more suppressed as the EC power is injected earlier.
- The  $T_e$  perturbation decreases asymmetrically in the radial direction. The magnetic island looks to decay from the smaller radius position.




# Dependence on injection timing



- When the ECCD is applied before the onset of NTM, the  $T_e$  perturbation is more suppressed.
- The critical EC timing for effective ECCD is associated with the NTM onset phase.



- The early ECCD is also effective in a high performance plasma of  $\beta_p \sim 1.7$ ,  $H_{89PL} \sim 1.8$ ,  $\beta_N \sim 3.0$  and  $H_{89PL} \beta_N \geq 5$ .
- The NB power for keeping  $\beta_N$  is lower when the 3/2 NTM is suppressed.
- The stabilization effect is sensitive to the EC current position as well as the fundamental early ECCD.



$I_p = 0.85$  MA,  
 $B_t = 1.7$  T,  $q_{95} = 3.5$   
 $P_{EC} = 2$  MW  
 $J_{EC} = 0.3 - 0.4$  MW/m<sup>2</sup>  
 $\rho_{EC} = 0.42 - 0.50$





## Modified Rutherford equation

$$\frac{\mu_0}{\eta} \frac{dW}{dt} = \underbrace{k_c \Delta'(W) \langle |\nabla \rho|^2 \rangle}_{\text{Classical}} + \underbrace{k_{BS} \mu_0 L_q j_{BS} \left\langle \frac{|\nabla \rho|}{B_p} \right\rangle \frac{W}{W^2 + W_d^2}}_{\text{Bootstrap}} - \underbrace{k_{GGJ} \varepsilon_s^2 \beta_p \frac{L_q^2}{\rho_s L_p} \left(1 - \frac{1}{q^2}\right) \langle |\nabla \rho|^2 \rangle \frac{1}{W}}_{\text{GGJ}}$$

$$- \underbrace{k_{pol} \varepsilon_s^{1.5} \beta_p \left( \frac{\rho_{pi} L_q}{L_p} \right)^2 \langle |\nabla \rho|^2 \rangle \frac{1}{W^3}}_{\text{Polarization}} - \underbrace{k_{EC} \mu_0 \frac{L_q}{\rho_s} \left\langle \frac{|\nabla \rho|}{B_p} \right\rangle \eta_{EC} \frac{I_{EC}}{a^2} \frac{1}{W^2}}_{\text{ECCD}}$$

ECCD term

EC code

Physical values  
at rational surface

Current profile

Additional  
current source

Geometry &  
Plasma profiles

## 1.5D tokamak simulation code ( TOPICS )

1D transport equations for density and temperatures

1D current diffusion equation

2D MHD equilibrium: Grad-Shafranov equation ( Fixed boundary )

Parameters of  $k_{BS}$ ,  $k_{GGJ}$ ,  $k_{pol}$ ,  $k_{EC}$  are constant.


Value of  $W_d$  depends on theoretical models to limit the parallel heat transport ).

( $k_{BS}$ ,  $k_{GGJ}$ ,  $k_{pol}$ ,  $k_{EC}$ ,  $W_d$ ) should be estimated by fitting to experiments.

# Parameter fitting at NTM growing and stabilizing phases



## Growth phase

$k_c = 1.2$ ,  $k_{BS} \sim 5$ ,  
 $k_{GGJ} < 10$  ( not important ),  
 $k_{pol} < 3$ ,  
 $W_d = 0.008$   
 ( ~ flux-limit model )



## Stabilization phase

$k_c = 1.2$ ,  $k_{BS} = 4.5$ ,  
 $k_{GGJ} = 1$ ,  $k_{pol} = 1$ ,  
 $W_d = 0.02$   
 ( ~ flux-limit model )  
 $k_{EC} \sim 3-4$ ,  $\Delta\rho_{EC} = 0.025$




Estimated coefficients are consistent between the growing and stabilizing phases.


# The rational surface is moved by EC current, resulting that the stabilizing effect is reduced.

- Higher EC current induces larger movement of the 3/2 surface.


$$I_{EC} / I_p = 0.02$$



$$I_{EC} / I_p = 0.04$$



- The power dependence in late ECCD agrees with the simulation.
- The EC power reduction for complete stabilization can be explained by the simulation. The NTM starts to grow with larger seed island width.



- The decrease of the bootstrap current term and/or the increase of classical tearing parameter term may contribute to the smaller island width in early ECCD.

	Bootstrap current term (Destabilizing term)	Classical tearing term (Stabilizing term)
$P_{\text{EC}}=0.8$ MW ( $I_{\text{EC}}=20$ kA)	-37%	+117%
$P_{\text{EC}}=1.7$ MW ( $I_{\text{EC}}=43$ kA)	-23%	+71%

- The stabilization of  $m=3/n=2$  NTM has been studied experimentally and theoretically in high  $\beta_p$  ELMy H-mode plasmas on the JT-60U tokamak.
- The early ECCD is more effective than the late ECCD, and the critical EC timing is associated with the mode onset phase.
- The  $3/2$  NTM has been suppressed in a high  $\beta_p$  ELMy H-mode plasma of  $\beta_N=3.0$  by the second harmonic early ECCD. The dependence on the EC current position is similar to the fundamental ECCD stabilization.
- The simulation on the basis of the modified Rutherford equation with 1.5D transport code and EC code well reproduce the time evolution of the  $3/2$  NTM experimentally observed both at the growing and stabilizing phases.
- The mode evolution can be explained by the simulation in late ECCD case. The EC power reduction for complete stabilization in early ECCD agrees with the simulation result.